## 7 MILLION DEATHS ANNUALLY LINKED TO AIR POLLUTION

GENEVA, 25 March 2014 – In new estimates released today, the World Health Organization (WHO) reports that in 2012 around 7 million people died – one in eight of total global deaths – as a result of air pollution exposure. This finding more than doubles previous estimates and confirms that air pollution is now the world's largest single environmental health risk. Reducing air pollution could save millions of lives.

In particular, the new data reveal a stronger link between both indoor and outdoor air pollution exposure and cardiovascular diseases, such as strokes and ischaemic heart disease as well as between air pollution and cancer. This is in addition to air pollution's role in the development of respiratory diseases, including acute respiratory infections and chronic obstructive pulmonary diseases

The new estimates are not only based on more knowledge about the diseases caused by air pollution, but also upon better assessment of human exposure to air pollutants through the use of improved measurements and technology. This has enabled scientists to make a more detailed analysis of health risks from a wider demographic spread that now includes rural as well as urban areas.

Regionally, low- and middle-income countries in the WHO South-East Asia and Western Pacific Regions had the largest air pollution-related burden in 2012, with a total of 3.3 million deaths linked to indoor air pollution and 2.6 million deaths related to outdoor air pollution.

"Cleaning up the air we breathe prevents noncommunicable diseases as well as reduces disease risks among women and vulnerable groups, including children and the elderly," says Dr Flavia Bustreo, WHO Assistant Director-General Family, Women and Children's Health. "Poor women and children pay a heavy price from indoor air pollution since they spend more time at home breathing in smoke and soot from leaky coal and wood cook stoves."

Included in the assessment is a breakdown of deaths attributed to specific diseases, underlining that the vast majority of air pollution deaths are due to cardiovascular diseases as follows:

Outdoor air pollution-caused deaths - breakdown by disease:

- 40% ischaemic heart disease;
- 40% stroke;
- 11% chronic obstructive pulmonary disease (COPD);
- 6% lung cancer;
- 3% acute lower respiratory infections in children.

Indoor air pollution-caused deaths – breakdown by disease:

- 34% stroke;
- 26% ischaemic heart disease;
- 22% COPD;
- 12% acute lower respiratory infections in children;
- 6% lung cancer.

The new estimates are based on the latest WHO mortality data from 2012 as well as evidence of health risks from air pollution exposures. Estimates of people's exposure to outdoor air pollution in different parts of the world were formulated through a new global data mapping. This incorporated satellite data, ground-level monitoring measurements and data on pollution emissions from key sources as well as modelling of how pollution drifts in the air.

"The risks from air pollution are now far greater than previously thought or understood, particularly for heart disease and strokes," says Dr Maria Neira, Director of WHO's Department for Public Health, Environmental and Social Determinants of Health. "Few risks have a greater impact on global health today than air pollution; the evidence signals the need for concerted action to clean up the air we all breathe."

After analysing the risk factors and taking into account revisions in methodology, WHO estimates indoor air pollution was linked to 4.3 million deaths in 2012 in households cooking over coal, wood and biomass stoves. The new estimate is explained by better information about pollution exposures among the estimated 2.9 billion people living in homes using wood, coal or dung as their primary cooking fuel, as well as evidence about air pollution's role in the development of cardiovascular and respiratory diseases, and cancers.

Contd. on page 59

## 7 MILLION DEATHS ANNUALLY LINKED TO AIR POLLUTION

Cont. from page 53

After analysing the risk factors and taking into account revisions in methodology, WHO estimates indoor air pollution was linked to 4.3 million deaths in 2012 in households cooking over coal, wood and biomass stoves. The new estimate is explained by better information about pollution exposures among the estimated 2.9 billion people living in homes using wood, coal or dung as their primary cooking fuel, as well as evidence about air pollution's role in the development of cardiovascular and respiratory diseases, and cancers.

In the case of outdoor air pollution, WHO estimates there were 3.7 million deaths in 2012 from urban and rural sources worldwide.

Many people are exposed to both indoor and outdoor air pollution. Due to this overlap, mortality attributed to the two sources cannot simply be added together, hence the total estimate of around 7 million deaths in 2012.

"Excessive air pollution is often a by-product of unsustainable policies in sectors such as transport, energy, waste management and industry. In most cases, healthier strategies will also be more economical in the long term due to health-care cost savings as well as climate gains," says Dr Carlos Dora, WHO Coordinator for Public Health, Environmental and Social Determinants of Health. "WHO and health sectors have a unique role in translating scientific evidence on air pollution into policies that can deliver impact and improvements that will save lives."

The release of today's data is a significant step in advancing a WHO roadmap for preventing diseases related to air pollution. This involves the development of a WHO-hosted global platform on air quality and health to generate better data on air pollution-related diseases and strengthened support to countries and cities through guidance, information and evidence about health gains from key interventions.

Later this year, WHO will release indoor air quality guidelines on household fuel combustion, as well as country data on outdoor and indoor air pollution exposures and related mortality, plus an update of air quality measurements in 1600 cities from all regions of the world.

World Health Organization. Media centre. 7 million premature deaths annually linked to air pollution [Internet]. Geneva: WHO; 2014 [cited 2014 Mar 27]. Available from: http://www.who.int/mediacentre/news/releases/2014/air-pollution/en/.